

# Tweed Shire Community Greenhouse Gas Profile Report



Prepared for  
Tweed Shire Council

Version	Author	Date	Description of changes
V0a - b	Nick Cody	30/10/2017	First draft
V1a	Hannah Snape	10/11/2017	Final draft
V1b - d	Nick Cody	14/02/2018	Final report after feedback from Council

Prepared by  
Ironbark Sustainability  
Suite 8, 70-80 Wellington St, Collingwood 3066  
ABN: 51 127 566 090  
Ph: 1300 288 262 | [info@realaction.com.au](mailto:info@realaction.com.au) | [www.realaction.com.au](http://www.realaction.com.au)

© 2017 Ironbark Group Pty. Ltd.

The information contained in this document produced by Ironbark Group Pty. Ltd is solely for the use of the client identified on this page for the purpose for which it has been prepared and Ironbark Group Pty. Ironbark undertakes no duty to or accepts any responsibility to any third party who may rely upon this document. All rights reserved. No section or element of this document may be removed from this document, reproduced, electronically stored or transmitted in any form without the written permission of Ironbark Group Pty. Ltd.

#### About Ironbark Sustainability

Ironbark Sustainability is a specialist consultancy that works with government and business around Australia by assisting them to reduce energy and water usage through sustainable asset and data management and on-the-ground implementation.

Ironbark has been operating since 2005 and brings together a wealth of technical and financial analysis, maintenance and implementation experience in the areas of building energy and water efficiency, public lighting and data management. We pride ourselves on supporting our clients to achieve real action regarding the sustainable management of their operations.

#### Our Mission

The Ironbark mission is to achieve real action on sustainability for councils and their communities.

# Contents

Abbreviations and Definitions	5
1. Executive Summary	7
2. Introduction	9
2.1 Background.....	9
2.2 Council Community Profile .....	9
2.3 Use of this Profile.....	10
3. Methodology and GPC Compliance	11
3.1 Disclaimer.....	13
3.2 IPCC Greenhouse Gas System Boundary.....	13
4. Total Emissions Profile	15
4.1 Emissions Breakdown.....	15
4.2 Completeness.....	17
4.3 Data Accuracy .....	18
5. Recommendations and Next Steps	20
5.1 Proposed Next Steps .....	20
6. Appendix 1: Data Inputs	21
6.1 Source Data .....	21
6.2 Greenhouse Gas Emissions Factors .....	22
7. Appendix 2: Policy Context	27

## Tables

Table 1: Total community emissions summary for Tweed Shire Council .....	7
Table 2: Sector descriptions used for GPC profiles.....	11
Table 3: Total community emissions summary according to sector.....	15
Table 4: Data accuracy classifications.....	19
Table 5: Grid supplied electricity emissions factors (Source: National GHG Accounting Factors 2016).....	22
Table 6: Grid supplied gas emissions (Source: National GHG Accounting Factors 2015, Table 2).....	22
Table 7: Natural gas leakage factors (Source: National GHG Accounting Factors 2015, Table 16).....	22
Table 8: Distribution losses (Source: Electricity and Gas Australia 2015, Table 3.5) .....	23
Table 9: Waste emission factors (Source: National GHG Accounting Factors 2015).....	23
Table 10: Waste emission factors for total waste disposed to landfill by broad waste stream category (Source: National GHG Accounting Factors 2015) .....	23
Table 11: Transport fuels emission coefficients (Source: National GHG Accounting Factors 2015, Table 4) .....	24
Table 12: Emission breakdown by fuel type (Source: National GHG Accounting Factors 2015) .....	25

Table 13: Global warming potentials of greenhouse gases (Source: National GHG Accounting Factors 2015, Appendix 1) ..... 25

Table 14: Electricity generation fuel mixes by state (Source: Electricity and Gas Australia 2015, Table 2.6) ..... 26

## Figures

Figure 1: Tweed Shire Council Municipal Boundary ..... 9

Figure 2: Emissions scopes definitions ..... 13

Figure 3: Emissions scopes and boundaries ..... 14

Figure 4: **Tweed Shire’s emissions by source** ..... 15

Figure 5: **Tweed Shire’s emissions by gas type and source** ..... 16

Figure 6: **Tweed Shire’s community emissions by scope** ..... 17

Figure 7: Notation keys ..... 18

## Abbreviations and Definitions

Term	Definition
ABS	Australian Bureau of Statistics
ABARES	Australian Bureau of Agricultural and Resource Economics and Sciences (previously Australian Bureau of Agricultural and Resource Economics or ABARE)
Activity Data	Activity data is a quantitative measure of a level of activity that results in GHG emissions taking place during a given period of time (e.g., volume of gas used, kilometres driven, tonnes of solid waste sent to landfill, etc.)
AEC	Australian Energy Council
AFOLU	Agriculture, Forestry, and Other Land Use
C40	C40 Cities Climate Leadership Group
CCA	Climate Change Authority
CCP	Cities for Climate Protection
cCR	Carbon Climate Register
CH <sub>4</sub>	Methane
CSIRO	Commonwealth Scientific and Industrial Research Organisation
CO <sub>2</sub>	Carbon dioxide
CO <sub>2</sub> -e	Carbon dioxide equivalent. The universal unit of measurement to indicate the global warming potential (GWP) of each GHG, expressed in terms of the GWP of one unit of carbon dioxide. It is used to evaluate the climate impact of releasing (or avoiding releasing) different GHGs on a common basis.
Global Covenant of Mayors	A coalition of city leaders addressing climate change by pledging to reduce their greenhouse gas emissions, tracking their progress and preparing for the impacts of climate change.
DNSP	Distribution Network Service Provider (Electricity Network)
Emissions Factor	An emissions factor is a measure of the mass of GHG emissions relative to a unit of activity.
ERF	Emissions Reduction Fund
ENA	Electricity Networks Australia
GHG	Greenhouse Gas
GHG Protocol	The Greenhouse Gas Protocol, developed by World Resources Institute and World Business Council on Sustainable Development, sets the global standard for how to measure, manage, and report GHG emissions.
GPC	Global Protocol for Community-Scale Greenhouse Gas Emission Inventories
GRP	Gross Regional Product
GSP	Gross State Product
Gt	Gigatonne

GWP	Global Warming Potential. The Global Warming Potential was developed to allow comparisons of the global warming impacts of different gases. Specifically, it is a measure of how much energy the emissions of one tonne of a gas will absorb over a given period of time, relative to the emissions of one tonne of carbon dioxide
ICLEI	International Council for Local Government Initiatives
IPCC	Intergovernmental Panel on Climate Change
IPPU	Industrial Processes and Product Use
kt	Kilotonne
Mt	Megatonne
MCF	Methane Correction Factor
SBT	Science-Based Targets
SEIFA	Socio-Economic Indexes for Areas
WRI	World Resources Institute

# 1. Executive Summary

This report provides an overview of the community greenhouse gas (GHG) profile for Tweed Shire Council. As a signatory to the Global Covenant of Mayors for Climate & Energy (herein referred to as the Global Covenant of Mayors), Tweed Shire Council is required to develop an emissions profile compliant with the Global Protocol for Community-Scale Greenhouse Gas Emission Inventories (GPC). The development of this profile is the first step in the preparation of a full process for mitigating climate change.

In line with GPC requirements, this profile includes an emissions total, as well as exploring emissions through a number of filters: sector, gas type, and scope.

The total emissions summary is provided below (Table 1). **Council’s total annual community emissions** have been calculated as 880kt CO<sub>2</sub>-e. The largest source of emissions by sector in Tweed is stationary energy, which mainly comprises electricity consumed by buildings and facilities and accounts for 62% (541kt) of total emissions. Meanwhile, on-road transportation is responsible for 31% of emissions, solid waste accounts for 6% and emissions from wastewater 1%.

Table 1: Total community emissions summary for Tweed Shire Council

Category	Emissions (t CO <sub>2</sub> e)	Percentage (%)
Stationary Energy	541,419	62%
Transportation	269,613	31%
Waste	54,833	6%
Wastewater	11,551	1%
Total	877,416	100%

The majority of **Tweed’s community** emissions are scope 2, which refers to emissions that are released as a result of using grid electricity and gas within the municipality, but with the emissions from energy generation occurring **outside of Council’s boundary**. Emissions from energy use in residential, commercial and industrial sectors are the main sources of scope 2 emissions.

It is important to note that while this profile is compliant with the international standard (GPC), and covers emissions from the entire community, Council only has a limited amount of control and influence over many of the emissions sources. While Council can continue to advocate and assist the broader community to implement energy efficiency and renewable energy projects, a cross-sectoral approach – with residents, business and other levels of government – is required for substantial emissions reductions throughout the municipality.

Rather than use this profile as a baseline upon which to develop targets and measure the impact of climate change mitigation actions by Council, targeted monitoring should instead be used to measure and report the impact of these items. As data sources and emissions calculation methods improve into the future, these items will be **able to be linked to Council’s** profile as a monitoring tool.

Recommended next steps from here are:

- Work with other councils, greenhouse alliances, state government and stakeholders, such as ICLEI Oceania and Ironbark, to source improved and updated data that can feed into future profiles and improve accuracy.
- Continue to seek broader Global Covenant of Mayors compliance by developing community-wide emissions reduction targets. To be best-**practice this should be a “science-based” target** in line with maintaining global average temperatures below a 2-degree increase.
- Continue to undertake climate mitigation and adaptation actions (through the implementation of the Tweed Byron Climate Change Adaptation Plan and the North Coast Integrated Regional Vulnerability Assessment) and measure and report the impact of these actions through targeted monitoring.



## 2. Introduction

### 2.1 Background

Sourcing and analysis of community GHG emissions data has been a cause of frustration for Australian councils over a long period of time. A decade ago, hundreds of councils had access to “top-down” community data provided by ICLEI Oceania based on ABS (Australian Bureau of Statistics) and ABARES (Australian Bureau of Agricultural and Resource Economics and Sciences) data. However, the development of this data ceased in 2009 with the closure of the Cities for Climate Protection funding.



Over the last 12 months, a range of council stakeholders have been working on collating data from these traditional sources (ABS and ABARES) as well as directly from utilities, CSIRO and other state data (for example on waste and transport) to develop GPC-compliant community inventories.

This has included the development and submission of GPC-compliant community inventories for the purpose of complying with the Global Covenant of Mayors.

The development of the Tweed GHG profile has been undertaken by a team from Ironbark Sustainability who have extensive experience **working at and with councils**. **What’s more, members of the team have recently been certified through the World Bank’s City Climate Planner Certificate Program**, a training and professional certificate program that aims to increase the global talent base of climate planning professionals.

### 2.2 Council Community Profile

Tweed Shire Council is located in north eastern New South Wales. It shares its municipal boundary with Byron Shire Council and Lismore City Council in the south, Kyogle Shire Council in the west and the state border between New South Wales and Queensland in the north (Figure 1). To the east, 37kms of coastline runs north-south along the Pacific Ocean. The population of 2015/2016 GPC Profile year was estimated to be 96,034.

Tweed Shire Council is a biologically diverse region of NSW. The region has substantial development pressure from its close proximity to the major growth area of South East Queensland and the area has witnessed strong economic and population growth over the past decade. Council has measures to reduce impact of future urban development as well as climate change. These impacts are likely to affect biodiversity, water security, coastal retreat and property encroachment, flooding issues, and




Figure 1: Tweed Shire Council Municipal Boundary

impacts to agriculture and tourism business. In response to this, Council has developed a number of sustainability and greenhouse gas minimisation initiatives. In 2009, Council, in conjunction with Byron Shire Council, developed a Climate Change Adaptation Plan to investigate risks in the region and ways to mitigate negative environmental, social and economic impacts. **In addition, Tweed Shire Council is included in the state government's** North Coast Integrated Regional Vulnerability Assessment of 2014. The assessment covers the vulnerability of human services, landscapes and ecosystems, industries, settlements and infrastructure, and emergency management sectors to a changing climate. Both the Regional Vulnerability Assessment and the Climate Change Adaptation Plan point to the ongoing pressures of growth in the area requiring a strategic approach to managing economic and population growth while minimising greenhouse gas emissions and adapting to changing climatic conditions.

## 2.3 Use of this Profile

It is very important that this profile is read and used appropriately. It should be stressed that whilst this profile contains a snapshot of the GHGs allocated to activity within Tweed municipality, it is not currently suitable as a monitoring tool for measuring the success of targets or actions.

A GPC-compliant profile is required in order to comply with the requirements of the Global Covenant of Mayors, to which Tweed Shire Council has committed. The GPC values completeness over accuracy, which ensures that when profiles are compiled across municipalities, emissions are not double-counted nor unaccounted for. This means that it is sometimes preferable to use state or national level data scaled down to a municipal level, over more detailed data that may not be complete or correctly aligned to municipal boundaries.

This profile contains a mixture of modelled and detailed data. As new data sources become available and methods for calculating emissions become more sophisticated, the calculated profile for Tweed will change, regardless of actual changes to emissions.

Ironbark envisions that council targets and action plans will be able to link together with their GHG profiles in the future. Until this time, action plans and targets should still be developed and monitored separately from this profile.

### 3. Methodology and GPC Compliance

The GPC requires cities to measure and disclose a profile of GHG emissions and to total these emissions using two distinct but complementary approaches. One captures emissions from both production and consumption activities taking place within the city boundary, including some emissions released outside the city boundary. The other categorises **all emissions into “scopes,”** depending on where they physically occur. Separate accounting of emissions physically released within the city boundary should be used for aggregation of multiple city inventories in order to avoid double counting.

The GPC enables a municipality’s **emissions to be broken down into the five sectors** (where relevant data is available), summarised in Table 2.

Table 2: Sector descriptions used for GPC profiles

Sector	Description
Stationary Energy	Stationary energy sources are one of the largest contributors to a <b>city’s GHG emissions. These emissions come from the combustion of</b> fuel in residential, commercial and institutional buildings and facilities, and manufacturing industries and construction, as well as power plants to generate grid-supplied energy. This sector also includes fugitive emissions, which typically occur during extraction, transformation, and transportation of primary fossil fuels.
Transportation	Transportation covers all journeys by road, rail, water and air, including inter-city and international travel. GHG emissions are produced directly by the combustion of fuel or indirectly by the use of grid-supplied electricity. Collecting accurate data for transportation activities, calculating emissions and allocating these emissions to cities can be a particularly challenging process. To accommodate variations in data availability, existing transportation models, and profile purposes, the GPC offers additional flexibility in calculating emissions from transportation.
Waste (solid waste and wastewater)	Waste disposal and treatment produces GHG emissions through aerobic or anaerobic decomposition, or incineration. GHG emissions from solid waste are calculated by disposal route, namely landfill, biological treatment and incineration and open burning. If methane is recovered from solid waste or wastewater treatment facilities as an energy source, it is reported under Stationary Energy. Similarly, emissions from incineration with energy recovery are reported under Stationary Energy.
Industrial Processes and Product Use (IPPU)	GHG emissions are produced from a wide variety of non-energy related industrial activities. The main emission sources are releases from industrial processes that chemically or physically transform materials (e.g., the blast furnace in the iron and steel industry, and ammonia and other chemical products manufactured from fossil fuels and used as chemical feedstock).  During these processes, many different GHGs can be produced. In addition, certain products used by industry and end-consumers, such as refrigerants, foams or aerosol cans, also contain GHGs which can be released during use and disposal.

<p>Agriculture, Forestry and Other Land Use (AFO LU)</p>	<p>Emissions from the Agriculture, Forestry and Other Land Use (AFOLU) sector are produced through a variety of pathways, including livestock (enteric fermentation and manure management), land use and land use change (e.g., forested land being cleared for cropland or settlements), and aggregate sources and non-CO<sub>2</sub> emission sources on land (e.g., fertilizer application and rice cultivation). Given the highly variable nature of land-use and agricultural activity across geographies, GHG emissions from AFOLU are amongst the most complex categories for GHG accounting.</p>
----------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

The GPC provides overarching and sector-specific reporting guidance for sourcing data and calculating emissions. Councils should select the most appropriate methodologies based on the **purpose of their profile, availability of data, and consistency with their country’s national profile** and/or other measurement and reporting programs in which they participate. The GPC does not require specific methodologies to be used to produce emissions data; rather it specifies the principles and rules for compiling a city-wide GHG emissions profile. Where relevant, the GPC recommends using methodologies aligned with the 2006 IPCC Guidelines for National Greenhouse Gas Inventories.

Ironbark, in partnership with HuxConnect, has developed a GHG activity data tool that has been approved by ICLEI Oceania as being GPC-compliant.

**In compliance with the GPC’s BASIC levels of GHG emissions accounting, Council’s** community emissions profile includes scope 1, 2 and 3 emissions as required under the GPC for each sector (stationary energy, transportation and waste). Specifically, for GPC BASIC compliance, the inclusion of emissions for the following sources and scopes are required:

- All scope 1 emissions from stationary energy sources (excluding energy production supplied to the grid, which shall be reported in the scope 1 total)
- All scope 1 emissions from transportation sources
- All scope 1 emissions from waste sources (excluding emissions from imported waste, which shall be reported in the scope 1 total)
- All scope 2 emissions from stationary energy sources and transportation
- Scope 3 emissions from treatment of exported waste

Emissions were further defined by gas type as either carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>) or nitrous oxide (N<sub>2</sub>O) and then converted into carbon dioxide equivalent (CO<sub>2</sub>-e) for comparison and to measure total impact.

### 3.1 Disclaimer

This profile has been developed in accordance with the GHG Protocol Global Protocol for Community-Scale Greenhouse Gas Emission Inventories (GPC) by experts certified under the City Climate Planner Certificate Program.

The GPC is designed to promote best practice GHG accounting and reporting that was developed through an inclusive multi-stakeholder process. This involved input from experts from nongovernmental organisations, governments, and others convened by WRI, C40 and ICLEI.



Data has been sourced from a variety of third parties (such as electricity providers and the CSIRO). While every effort has been made to use data from reputable sources and a thorough quality assurance process undertaken, neither Council nor Ironbark Sustainability are responsible for data inaccuracies by third parties.

### 3.2 IPCC Greenhouse Gas System Boundary

To comply with the GPC, and indeed for any best-practice GHG profile development, a council (or organisation or sub-national body) must first define a profile boundary. This identifies the geographic area, time span, gases, and emission sources, covered by a GHG profile. Any geographic boundary may be used for the GHG profile.

For this profile, the boundary aligns with the administrative boundary of Tweed Shire and accounts for the GHG emissions in the reporting year 2015-2016. It covers the seven gases covered by the Kyoto Protocol.

Activities taking place within Tweed can generate GHG emissions that occur inside the city boundary as well as outside the city boundary. To distinguish between them, the GPC groups emissions into three categories based on where they occur: scope 1, scope 2 or scope 3 emissions. Definitions are provided in Figure 2, based on an adapted application of the scopes framework used in other international GHG protocol and standards.

Scope	Definition
<b>Scope 1</b>	GHG emissions from sources located within the city boundary
<b>Scope 2</b>	GHG emissions occurring as a consequence of the use of grid-supplied electricity, heat, steam and/or cooling within the city boundary
<b>Scope 3</b>	All other GHG emissions that occur outside the city boundary as a result of activities taking place within the city boundary

Figure 2: Emissions scopes definitions

Councils, by virtue of their size and connectivity, inevitably give rise to GHG emissions beyond their boundaries. Measuring these emissions allows cities to take a more holistic approach to tackling climate change by assessing the GHG impact of their supply chains, and identifying areas of shared responsibility for upstream and downstream GHG emissions.

The GPC also includes scope 3 accounting for a limited number of emission sources, including transmission and distribution losses associated with grid-supplied energy, and waste disposal and treatment outside the city boundary and transboundary transportation.

The scopes framework helps to differentiate emissions occurring physically within the city (scope 1), from those occurring outside the city (scope 3) and from the use of electricity, steam, and/or heating/cooling supplied by grids which may or may not cross city boundaries (scope 2). **Scope 1 emissions may also be termed "territorial" emissions because they occur discretely within the territory defined by the geographic boundary.**

Figure 3 illustrates which emission sources occur solely within the geographic boundary established for the profile, which occur outside the geographic boundary, and which may occur across the geographic boundary.




Figure 3: Emissions scopes and boundaries

## 4. Total Emissions Profile

### 4.1 Emissions Breakdown

Council’s total annual community emissions have been calculated as 880kt CO<sub>2</sub>-e.

Table 3: Total community emissions summary according to sector

Category	Emissions (t CO <sub>2</sub> e)	Percentage (%)
Stationary Energy	541,419	62%
Transportation	269,613	31%
Waste	54,833	6%
Wastewater	11,551	1%
Total	877,416	100%

As seen in Table 3 and Figure 4, the largest source of total GHG emissions is stationary energy, which includes residential buildings; commercial and institutional facilities; and manufacturing and construction industries.

On-road transportation produced the next highest emissions per category, 31% total emissions or 270kt CO<sub>2</sub>-e. No emissions have been included under transport sector for rail, aviation, water transport or other off-road transportation sources as this data was not readily available. Where transport sources are responsible for emissions through use of electricity, such as rail, these emissions have been included under stationary energy. As more granular data becomes available the reporting of transport emissions will be improved upon.




Figure 4: Tweed Shire’s emissions by source

Waste sources produced 7% of total emissions at 69kt CO<sub>2</sub>-e. The sources of this waste were from the biological treatment of wastewater (1%) including sewerage, and also from the disposal of solid waste (6%).

Figure 5 compares the emissions by source type and also shows the amounts of each GHG that is produced either directly or indirectly through activities within the municipality. Gases are measured in carbon dioxide equivalent to ensure they are comparable. Both stationary energy and transportation are largely responsible for releasing carbon dioxide with trace amounts of methane and nitrous oxide, whilst emissions from waste are mostly released as methane (CH<sub>4</sub>).




Figure 5: **Tweed Shire's** emissions by gas type and source






Figure 6: Tweed Shire’s community emissions by scope

NB: In Figure 6 Stationary energy has been broken down into 3 subcategories: residential, commercial and manufacturing.

The majority of Tweed’s community emissions are scope 2, which means GHGs are being released into the atmosphere at the source of energy generation, outside of Council’s boundary, however the energy is consumed within the boundary of Tweed Shire (Figure 6). Stationary energy is largely responsible for scope 2 emissions, generally via the electricity grid. Solid waste disposal is considered scope 3 emissions, because waste generated within the municipal boundary is transported to be treated outside of the boundary. Wastewater treatment is considered scope 1 emissions because waste that is generated by Tweed’s community is treated within the municipal boundary.

## 4.2 Completeness

Data collection is an integral part of developing and updating a GHG profile. Data will likely come from a variety of sources and will vary in quality, format, and completeness. In many cases, it needs to be adapted for the purposes of the profile. The GPC recognizes these challenges and sets out data collection principles and approaches. It also provides guidance on gathering existing data, generating new data and adapting data for profile use.

According to the GPC, a council’s GHG profile shall follow the principles of relevance, completeness, consistency, transparency and accuracy. In regards to completeness, a council should account for all required emissions sources within the profile boundary and any exclusion of emission sources shall be justified and clearly explained.

To accommodate for limitations in data availability and differences in emission sources between cities, the GPC requires the use of “notation keys”, as recommended in IPCC Guidelines. These are provided in Figure 7.

Notation key	Definition	Explanation
IE	Included Elsewhere	GHG emissions for this activity are estimated and presented in another category of the inventory. That category shall be noted in the explanation.
NE	Not Estimated	Emissions occur but have not been estimated or reported; justification for exclusion shall be noted in the explanation.
NO	Not Occurring	An activity or process does not occur or exist within the city.
C	Confidential	GHG emissions which could lead to the disclosure of confidential information and can therefore not be reported.

Figure 7: Notation keys

Where notation keys are used for this profile, an accompanying explanation is provided to justify exclusions or partial accounting of GHG emission source categories.

The GPC gives councils the option of selecting between two reporting levels: BASIC or BASIC+. The BASIC level covers emission sources that occur in almost all councils (stationary energy, in-boundary transportation, and in-boundary generated waste) and the calculation methodologies and data are more readily available. The BASIC+ level has a more comprehensive coverage of emissions sources (BASIC sources plus IPPU, AFOLU, transboundary transportation, and energy transmission and distribution losses) and reflects more challenging data collection and calculation procedures.

This profile satisfies the requirements of GPC BASIC. Localised granular data has been sought where possible, however time spent seeking granular data and/or BASIC+ level data has been capped.

### 4.3 Data Accuracy

Due to variances in the availability of some data there was the need to extrapolate and rely on assumptions in some cases.

The quality of data has been categorised and is outlined in Table 6. Further information on the classifications is provided below.

High - Detailed Activity Data: denotes data that was directly available from a reliable source, such as energy billing data provided by Council.

Medium - Modelled activity data using robust assumptions: where more reliable data is not available granular data has been modelled.

Low - Highly modelled or uncertain activity data: denotes data that was modelled from a highly reliable source, such as the Australian Bureau of Statistics.

Overall, data accuracy for this emissions profile is considered to be medium, but compliant with the GPC BASIC requirements. Recommendations for improving data accuracy are at Section 5.

Table 4: Data accuracy classifications

Methods			
Profile Key Parameters	Method Used	Data Reliability	
		Activity data	Emissions factors
Basic Municipal			
BM1	BM1.1	Detailed Activity Data	n/a
BM2	BM2.1	Detailed Activity Data	n/a
Stationary Energy			
SE1	SE1.1	Modeled activity data using robust assumptions	More general emission factors
SE2	SE2.1	Modeled activity data using robust assumptions	More general emission factors
SE3	SE3.1	Detailed Activity Data	More general emission factors
Transport			
TR1	TR1.1	Highly-modeled or uncertain activity data	More general emission factors
Waste			
WS1	WS1.1	Modeled activity data using robust assumptions	Default emission factors
WS2	WS2.3	Detailed Activity Data	Default emission factors

## 5. Recommendations and Next Steps

The development of this GPC-compliant profile is the first stage of compliance with the Global Covenant of Mayors, and should be considered part of a process towards a climate mitigation system involving targets and actions. This profile has been developed in a way that future improvements in data (be that around relevance, completeness, consistency, transparency and accuracy) can be easily adapted. This starting point is about developing a platform; reaching GPC BASIC compliance; ensuring there is no double-counting across council boundaries and that metrics are consistent across time.

A BASIC profile will only provide Global Covenant of Mayors compliance for one year, after which time the profile must be updated to BASIC+. This update will need to be of a higher standard, for example by incorporating more GHGs or more sectors.

As discussed in Section 0, this profile should not be used as a baseline for developing and monitoring targets and actions. Instead, it should provide a summary of the relative GHGs for which Tweed Shire is responsible and sit alongside, rather than above, these other items.

In the meantime, it is recommended that Council maintain their compliance with Global Covenant of Mayors through the development of a science-based target and implementation of their current climate action plan. Despite our recommendation to de-couple these items from a GHG profile for now, the savings from each action can still be closely monitored and reported separately to this profile.

It is expected that there will be changes to this profile as more granular data becomes available and methods for calculating and allocating emissions become more sophisticated. In the future, **Council's** community GHG profile will be considered accurate enough to link with their targets and actions as a full climate change mitigation system.

### 5.1 Proposed Next Steps

From here there are a number of actions that Council may undertake to maintain compliance with Global Covenant of Mayors and work towards a full climate mitigation system:

- Work together with other councils, greenhouse alliances, state government or organisations, such as Ironbark, to source improved and updated data that can feed into future profiles.
- Continue to seek broader Global Covenant of Mayors compliance by developing community-wide emissions reduction targets. To be best-practice this should be a **"science-based" target** in line with keeping global average temperatures below a 2-degree increase.
- Continue to undertake climate mitigation and adaptation actions (through the implementation of the Tweed Byron Climate Change Adaptation Plan and the North Coast Integrated Regional Vulnerability Assessment, and future iterations) and measure and report the impact of these actions through targeted monitoring.

## 6. Appendix 1: Data Inputs

### 6.1 Source Data

The following sources have been used to develop this profile.

NB: While Ironbark Sustainability has attempted to retrieve sources published at dates commensurate to the activity data generated in the community emissions profile, due to a number of factors around data confidence and publishing cycles, the quality of the reporting would be diminished if reports and recommendations are based on data from poor sources. Ironbark is always assessing the availability of data sources and aims to use more up to date and better quality sources where possible.

Name	Author	Publication date
Electricity Gas Australia 2015	Electricity Gas Australia	April 2015
Small-scale Technology Certificates - Registered	Clean Energy Regulator	April 2015
Population Estimates by Local Government Area 2005 to 2015	Australian Bureau of Statistics	September 2016
National Postcode Concordances 2011	Australian Bureau of Statistics	March 2016
National Regional Profile - Industry - LGA 2010-14	Australian Bureau of Statistics	June 2012
Waste Generation and Resource Recovery – 2010/11	Randell Environmental Consulting	February 2014
Waste Account, Australia, Experimental Estimates	Australian Bureau of Statistics	February 2013
National Greenhouse Inventory 2015	Australian Department of Environment and Energy	June 2016
National Greenhouse Accounting Factors	Department of Environment and Energy	July 2017
National Greenhouse Accounting Factors	Department of Environment and Energy	August 2015
National Greenhouse Inventory 2015	Department of Industry, Innovation and Science	October 2016
Australian Energy Statistics - 2016 Update	Electricity Gas Australia 2015	April 2015
Electricity Gas Australia 2015	Department of Industry, Innovation and Science	October 2016
National Greenhouse Gas Inventory	Australian Department of Environment and Energy	June 2014

## 6.2 Greenhouse Gas Emissions Factors

Emission factors convert activity data into a mass of GHG emissions; tonnes of CO<sub>2</sub> released per kilometre travelled, for example, or the ratio of CH<sub>4</sub> emissions produced to amount of waste landfilled. According to the GPC, emission factors should be relevant to the profile boundary, specific to the activity being measured, and sourced from credible government, industry, or academic sources<sup>1</sup>. The following tables outline the GHG emission factors used in the development of this profile.

Table 5: Grid supplied electricity emissions factors (Source: National GHG Accounting Factors 2016)

Grid supplied electricity emissions factors				
State	Financial Year	Equivalent year	Scope 2 EF (kg CO <sub>2</sub> -e/kWh)	Scope 3 EF (kg CO <sub>2</sub> -e/kWh)
NSW	Latest Estimate	2015.5	0.83	0.12

Table 6: Grid supplied gas emissions (Source: National GHG Accounting Factors 2015, Table 2)

Emission factors for the consumption of natural gas					
Fuel combusted	Energy content factor	Emission factor (kg CO <sub>2</sub> -e/GJ)			
	(GJ/m <sup>3</sup> unless otherwise indicated)	(relevant oxidation factors incorporated)			
		CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	Total
Natural gas distributed in a pipeline	39.3 × 10 <sup>-3</sup>	51.4	0.1	0.03	51.53

Table 7: Natural gas leakage factors (Source: National GHG Accounting Factors 2015, Table 16)

Natural gas leakage factors			
State	Unaccounted for gas (%UAG)	Natural gas composition factor (tonnes CO <sub>2</sub> -e/TJ)	
	UAG	CO <sub>2</sub>	CH <sub>4</sub>
NSW	2.2	0.8	390

<sup>1</sup> If no local, regional, or country-specific sources are available, councils can use IPCC default factors or data from the Emission Factor Database (EFDB) or other standard values from international bodies that reflect national circumstances.

Table 8: Distribution losses (Source: Electricity and Gas Australia 2015, Table 3.5)

Distribution losses	
Data year	2013.5
State	NSW
Loss	4.5%

Table 9: Waste emission factors (Source: National GHG Accounting Factors 2015)

Waste – Emissions Factors				
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	Total CO <sub>2</sub> -e
Proportion of total CO <sub>2</sub> -e	0.27%	96.08%	3.65%	100%

Table 10: Waste emission factors for total waste disposed to landfill by broad waste stream category (Source: National GHG Accounting Factors 2015)

Waste emission factors for total waste disposed to landfill by broad waste stream category			
Waste types	Municipal solid waste	Commercial and industrial waste	Construction / demolition waste
	A	B	C
Emission factor (t CO <sub>2</sub> -e/t waste)	1.4	1.3	0.2

Table 11: Transport fuels emission coefficients (Source: National GHG Accounting Factors 2015, Table 4)

Fuel combustion emission factors - fuels used for transport energy purposes				
Fuel combusted	Energy content factor (GJ/kL unless otherwise indicated)	Emission factor (kg CO <sub>2</sub> -e/GJ) (relevant oxidation factors incorporated)		
		CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
Pre-2004 vehicles				
Gasoline (other than for use as fuel in an aircraft)	34.2	67.4	0.5	1.8
Diesel oil	38.6	69.9	0.1	0.5
Liquefied petroleum gas	26.2	60.2	0.6	0.7
Ethanol for use as fuel in an internal combustion engine	23.4	0	0.7	1.9
Post-2004 vehicles				
Gasoline (other than for use as fuel in an aircraft)	34.2	67.4	0.02	0.2
Diesel oil	38.6	69.9	0.01	0.6
Liquefied petroleum gas	26.2	60.2	0.4	0.3
Ethanol for use as fuel in an internal combustion engine	23.4	0	0.2	0.2



Table 12: Emission breakdown by fuel type (Source: National GHG Accounting Factors 2015)

Emissions breakdown by fuel type							
Data year	2013.5						
Fuel type	CO <sub>2</sub> emissions factors (kg CO <sub>2</sub> -e/GJ)				Proportion of emissions		
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	Total	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
Hydro	0.00	0.00	0.00	0.00	0.0000	0.0000	0.0000
Biofuels	0	0.07	1.1	1.17	0.0000	0.0598	0.9402
Black coal	90	0.03	0.2	90.23	0.9975	0.0003	0.0022
Brown coal	93.50	0.02	0.40	93.92	0.9955	0.0002	0.0043
Coal seam methane	51.4	0.2	0.03	51.63	0.9955	0.0039	0.0006
Natural gas	51.40	0.10	0.03	51.53	0.9975	0.0019	0.0006
Oil products	69.8	0	0.2	70.00	0.9971	0.0000	0.0029
Solar	0	0	0	0.00	0.0000	0.0000	0.0000
Wind	0.00	0.00	0.00	0.00	0.0000	0.0000	0.0000
Total (kg CO <sub>2</sub> -e/GJ)	81.08	0.02	0.35	81.46	0.9954	0.0003	0.0043
Total (kg CO <sub>2</sub> -e/MWh)	22.52	0.01	0.10	22.63	0.9954	0.0003	0.0043
	99.536%	0.029%	0.434%	1.00			

Table 13: Global warming potentials of greenhouse gases (Source: National GHG Accounting Factors 2015, Appendix 1)

Global Warming Potentials		
Gas	Chemical formula	Global Warming Potential
Carbon dioxide	CO <sub>2</sub>	1
Methane	CH <sub>4</sub>	25
Nitrous oxide	N <sub>2</sub> O	298

Table 14: Electricity generation fuel mixes by state (Source: Electricity and Gas Australia 2015, Table 2.6)

Electricity generation fuel mixes		
Data year	2013.5	
State	NSW	
Generation source	Value (GWh)	Proportion
Hydro	4,213.1	6.3%
Biofuels	882.3	1.3%
Black coal	54,626.3	81.1%
Brown coal	0	0.0%
Coal seam methane	0	0.0%
Natural gas	5,528.3	8.2%
Oil products	170.7	0.3%
Solar	1,067.4	1.6%
Wind	899.2	1.3%
Total (kg CO <sub>2</sub> -e/GJ)	67,387.3	100.0%
Fossil Fuels	60,325.3	90%
Nuclear	0	0%
Renewables	7,062.0	10%

## 7. Appendix 2: Policy Context

### International/Federal

Globally, a number of countries, states and cities are making significant changes to the way they live and work to reduce their contribution to human induced climate change. In some countries, such as Australia, it has been cities and communities that are leading the way with policies and programs to seize the opportunity of sustainability.

Australia has seen significant policy uncertainty through the introduction and repeal of policies such as the price on carbon and the protracted review of the Renewable Energy Target. Nationally, uncertainty regarding the Renewable Energy Target and carbon pricing continues to stifle investment in large-scale renewable energy to the detriment of consumer prices.

The second major Federal policy influencing Council decision-making is the Emission Reduction Fund (ERF). The methodologies and abatement pricing structures announced in the first round of the Fund provide limited opportunities for councils to participate, however the second round may provide some limited opportunity. The Australian Renewable Energy Agency and Clean Energy Finance Corporation are still attempting to fund innovation and tailored project finance to support the development of the industry irrespective of limited political support.

Australia recently became a signatory to the Paris Climate Agreement which pledges to reduce **carbon emissions and limit global warming to “well below 2°C above pre-industrial levels”**.

### State

The NSW Government has released the NSW Climate Change Policy Framework, which commits NSW to the aspirational objectives of achieving net-zero emissions by 2050 and helping NSW to become more resilient to a changing climate.

#### *Climate Change Fund*

The Climate Change Fund was established in 2007 to address the impacts of climate change, encourage energy and water saving activities and increase public awareness and acceptance of climate change.

On 3 November 2016, the NSW Government released a Draft Climate Change Fund Strategic Plan and A Draft Plan to Save NSW Energy and Money. The government ran public consultation from 3 November to 16 December 2016. The Draft Strategic Plan sets out priority investment areas and potential actions for up to \$500 million of new funding between 2017–18 and 2021–22.

#### *Energy Efficiency Action Plan*

The Energy Efficiency Action Plan (EEAP), launched in response to Goal 5 of ‘NSW 2021’, consists of 30 actions to strengthen the energy efficiency market and help NSW households, business and government to use energy more efficiently.

With the overall goal of tackling the costs of living in NSW, the EEAP aims to reach the following targets:

- achieve 16,000 GWh in energy savings per year by 2020
- support 220,000 low income households to reduce energy use by up to 20% by 2014
- assist 50% of NSW commercial floor space achieve a four-star NABERS energy and water rating by 2020, through the delivery of high-standard building retrofit programs

#### *Community Renewable Energy*

Through the Regional Clean Energy Program (RCEP), the NSW Government is helping communities to produce their own electricity locally, using renewable energy resources.

The RCEP is a key component of the NSW Renewable Energy Action Plan (REAP) to increase generation in the state, and work closely with communities and industry to bring new jobs, investment and technological advances to local economies.

#### Local

At the local level, a number of councils across Australia have shown strong leadership in climate action and derived great benefits by reducing their operational costs, carbon exposure and improving productivity. In addition, this has provided a great basis for community engagement and leadership.